

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



PROPYZAMIDE 500 SC

Herbicide

ACTIVE CONSTITUENT: 500 g/L PROPYZAMIDE

GROUP 3 HERBICIDE

For Selective Control of Certain Grasses and Broadleaf Weeds in Lettuce, Sports Turf, Home Lawns and Legumes Seed Crops and Pastures, as per the Direction for Use

IMPORTANT: READ THE ATTACHED BACK LABEL BEFORE OPENING OR USING THIS PRODUCT

Swan Chemical Holdings Pty. Ltd.

Address: u2/9 Glossop St Wangara WA 6065.

PH: 1300 289 520 E: info@swanchemicalholdings.com

W: swanchemicalholdings.com

20L

110L

1000L

SWAN

PROPYZAMIDE 500 SC HERBICIDE

DIRECTIONS FOR USE

RESTRAINTS

Spray drift restraints

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings 1 to 2 hours before sunset and persist until 1 to 2 hours after sunrise.

SITUATION	WEEDS CONTROLLED	STATE	RATE L/ha	CRITICAL COMMENTS	
Lettuce Direct sown Transplanted	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Ryegrass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles	All States	4.5 in 330 – 550L	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
				Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
		Vic only	2.3 in 300 – 550L	Apply on light sandy soils, low in organic matter.	
Turf & Lawns Common Couch Queensland Blue couch Buffalo grass	Winter grass				
				Rate Bowling Green 37 x 37m	
		All States	1.2 (12mL/100m ²)	170 mL	Apply during the autumn and Winter, if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12-25 mm irrigation or rain and again 24 hours later. Repeat if necessary on Qld. Blue Couch after 4 weeks. DO NOT re-seed for 60 days after treatment. Use ONLY on established turf. DO NOT use on Bluegrass, ryegrass, fescue or bent grass lawns.
		Qld only	0.6 (6mL/100m ²)	85 mL	
WA only	1.2 (12mL/100m ²)	170 mL			
Lucerne, Clover, Medics etc. grown for forage, hay or seed production	Annual Ryegrass, Winter grass, Barley grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue and Rats Tail Fescue Wild Oats, Perennial Ryegrass, Bentgrass, English Couch, Yorkshire Fog Grass, Chickweed, seeding Sorrel, Blackberry Nightshade, Patterson's Curse, seedling Wireweed	NSW, ACT, SA, WA, Tas only	1-1.5	SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July. WEEDS: Apply when weeds are at post or pre-emergent stages.	
			1.5-2	BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatments is recommended.	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OF FOR SEED FOR 25 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

MIXING

Prior to pouring, shake container vigorously, then add the required quantity of Spalding Propyzamide 500 SC Herbicide to water in the spray vat while stirring or with agitators in motion. **DO NOT** mix with other pesticides.

RESISTANT WEEDS WARNING

GROUP	3	HERBICIDE
-------	----------	-----------

Swan Propyzamide 500 SC Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management Swan Propyzamide 500 SC Herbicide is a Group D herbicide. Some naturally occurring weed biotypes resistant to Swan Propyzamide 500 SC Herbicide and other amide herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly.

These resistant weeds will not be controlled by Swan Propyzamide 500 SC Herbicide or other amide herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Swan Chemical Holdings Pty. Ltd. accepts no liability for any losses that may result from failure of Swan Propyzamide 500 SC Herbicide to control resistant weeds. Advice as to strategies and alternative treatments that can be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or Swan Chemicals Holdings representative.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or watercourses with the chemical or used containers. **DO NOT** graze sheep on treated areas or feed to poultry or pigs.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping land or pastures.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. Protect from frost.
Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.
If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in accordance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID INSTRUCTIONS

If poisoning occurs contact a doctor or Poisons Information Centre, Telephone Australia 13 11 26; New Zealand 0800 764 766

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet.

CONDITIONS OF SALE

The use of Swan Propyzamide 500 SC Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Swan Chemical Holdings Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Swan Chemical Holdings Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

**IN EMERGENCY DIAL
000
POLICE OR FIRE BRIGADE**

**24HR EMERGENCY RESPONSE:
AU +61 1800 951 288
NZ +64 800 700 112**

APVMA Approval No: 95321/145285

DOM:

Batch No:

